

POISON

KEEP OUT OF REACH OF CHILDREN
FIRE AND EXPLOSION HAZARD
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

MULTI TABS

MULTI FUNCTION STABILISED POOL SANITISER

TABLETS



ACTIVE CONSTITUENTS:
900g/Kg available CHLORINE (Cl) present as
TRICHLORO ISOCYANURIC ACID
7.5g/Kg COPPER present as
COPPER SULPHATE PENTAHYDRATE

APVMA: 59923/1105

CONDITION OF SALE

FOCUS PRODUCTS P/L will not accept any responsibility howsoever it will arise and whether consequently or otherwise in connection with the supply or use of these goods other than their responsibility for the merchantable quality of the goods and such responsibilities mandatorily imposed by applicable statutes. To the extent allowed by such statutes the liability of FOCUS PRODUCTS P/L is limited to the cost of the product purchased.

NOT TO BE USED FOR ANY PURPOSE OR IN ANY MANNER CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.

DIRECTIONS FOR USE

POOL VOLUME (Litres)	Recommended Weekly dose
20,000	1
40,000	2
60,000	3

The dosage rates are only a guide to achieve the required levels of free chlorine.

REQUIRED LEVELS	MIN FREE CHLORINE (mg/L)	TOTAL ALKALINITY	pH RANGE (optimum)	STABILISER (mg/L)
Cold Water	2	60-200	7.2 to 7.6	30-50
Water Temp >26°C	3			

DO NOT exceed a concentration of 50mg/L of isocyanurate.

GENERAL DIRECTIONS

The correct level of free chlorine should be maintained for several hours prior to and during swimming activity. Chlorine levels and pH should be tested daily by use of a reliable test kit using fresh testing reagents. If pH falls below the required level add FOCUS pH INCREASE (dry alkali), if it rises above, add FOCUS DRY ACID or Focus Hydrochloric Acid. Dose pool in the evening when the pool is not in use.

Stabiliser levels MUST be tested fortnightly by your local pool shop or by use of a reliable test kit. Once stabiliser levels reach 50mg/L DO NOT use this product until stabiliser levels fall below that level. INSTEAD use an unstabilised chlorine product to maintain free chlorine levels. If 50mg/L of stabiliser is exceeded the ability of the chlorine to control algae and bacteria is reduced. UNDER NO CIRCUMSTANCES should stabiliser concentrations be allowed to exceed 100mg/L. If stabiliser concentration needs to be reduced consult your local pool shop. Always add tablets to a floating dispenser device or feeder designed for either calcium hypochlorite tablets or Trichloroisocyanuric tablets.

DO NOT USE IN FEEDERS DESIGNED FOR ANY OTHER CHEMICAL SYSTEMS.

CAUTION

NOT TO BE USED AS A FOOD CONTAINER. THIS PRODUCT IS NOT SUITABLE FOR SUPER-CHLORINATION.

SAFETY DIRECTIONS

Highly reactive oxidising chlorine compound. May cause fire or explosion or produce severe burns. Product will irritate the eyes, nose, throat and skin. Avoid contact with the eyes, skin and clothing. DO NOT inhale dust. DO NOT allow product to come into contact with other chemicals, especially acids, or with combustible materials such as paper, fabric, sawdust or kerosene. DO NOT allow it to get damp. Use clean containers for dispensing. DO NOT mix with other chemicals. DO NOT mix with different types of chlorinating chemicals. DO NOT add water to product - add the product to water, but in case of fire, drench with water. Store undercover in a dry, clean, well ventilated place away from sunlight. Wash hands after use. Harmful if swallowed.

FIRST AID

If poisoning occurs contact a doctor or Poisons Information Centre (phone 13 1126) If swallowed, DO NOT induce vomiting. Give a glass of water. If skin contact occurs, remove contaminated clothing and wash skin thoroughly. If in eyes, hold eyes open, flood with water for at least 15 minutes and see a doctor.

IN EMERGENCY DIAL 000 POLICE OR FIRE BRIGADE,

SPECIALIST



ADVICE IN EMERGENCY ONLY - DIAL 1 800 033 111 ALL HOURS AUSTRALIA WIDE

**HAZCHEM 1W
UN 2880
PGII**

**IN EMERGENCY DIAL 000 (NZ 111)
POLICE OR FIRE BRIGADE**



© 2021 Focus Products Pty Ltd

All rights reserved.

Unauthorised reproduction is prohibited by law.

